

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number:

0 574 048 A2

(12)

EUROPEAN PATENT APPLICATION(21) Application number: **93200691.9**

(51) Int. Cl.⁵: **C12N 15/38, C12P 21/08,
C12Q 1/68, C07K 13/00,
C07K 15/28, G01N 33/536,
G01N 33/68**

(22) Date of filing: **10.03.93**

The microorganism(s) has (have) been deposited with European Collection of Animal Cell Cultures (ECACC) under number(s) 93020413, 932020412 and 93020414.

(30) Priority: **13.03.92 EP 92200721**

(43) Date of publication of application:
15.12.93 Bulletin 93/50

(84) Designated Contracting States:
**AT BE CH DE DK ES FR GB GR IE IT LI LU MC
NL PT SE**

(71) Applicant: **AKZO N.V.**
Velperweg 76
NL-6824 BM Arnhem(NL)


(72) Inventor: **Middeldorp, J.M.**
Staringstraat 151
NL-5343 GD Oss(NL)

(74) Representative: **Hermans, Franciscus G.M. et al**
P.O. Box 20
NL-5340 BH Oss (NL)

(54) **Peptides and nucleic acid sequences related to Epstein Barr Virus.**

(57) The present invention relates to peptides immunochemically reactive with antibodies to the Epstein-Barr virus (EBV), comprising at least part of the VCA-p18 or VCA-p40 protein, encoded within the EBV open reading frames BFRF3 and BdRF1 respectively, or a functional variant thereof. The invention further relates to nucleic acid sequences encoding these peptides, monoclonal antibodies and anti-idiotypic antibodies. The invention also relates to recombinant vector molecules comprising a nucleic acid sequence according to the invention and host cells transformed or transfected with these vector molecules. The invention is further concerned with immunological reagents and methods for the detection of EBV or anti-EBV antibodies and a method for the amplification and detection of Epstein Barr viral nucleic acid.

FIGURE 1



The present invention relates to peptides immunochemically reactive with antibodies to the Epstein-Barr virus (EBV), nucleic acid sequences encoding these peptides, monoclonal antibodies against these peptides, cell lines capable of producing monoclonal antibodies and anti-idiotypic antibodies. The invention also relates to recombinant vector molecules comprising a nucleic acid sequence according to the invention and host cells transformed or transfected with these vector molecules. The invention is further concerned with immunological reagents and methods for the detection of EBV or anti-EBV antibodies and a method for the amplification and detection of Epstein Barr viral nucleic acid.

EBV is an ubiquitous human herpes virus that was first discovered in association with the African (endemic or e) form of Burkitt's lymphoma (BL). Subsequently the virus was also found associated with nasopharyngeal carcinoma (NPC) and was shown to be the causative agent of infectious mononucleosis (IM). Infection usually occurs during early childhood, generally resulting in a subclinical manifestation, occasionally with mild symptoms. Infection during adolescence or adulthood, however, can give rise to IM characterized by the presence of atypical lymphocytes in the periphery. The bulk of these lymphocytes are T lymphocytes; however, included in their number are a small population of B lymphocytes infected by EBV. The infection of B lymphocytes may also be accomplished in vitro. Such cells become transformed and proliferate indefinitely in culture and have been referred to as "immortalized", "latently infected" or "growth transformed". As far as is known, all individuals who become infected with EBV remain latently infected for life. This is reflected by the lifelong continuous presence of small numbers of EBV-genome positive transformed B-cells among the circulating peripheral blood lymphocytes and the continuous but periodic shedding of virus in the oropharynx.

In the vast majority of cases EBV infection results in a lymphoproliferative disease that may be temporarily debilitating, but is always benign and self-limiting. In certain immunosuppressed individuals, however, the result can be full-blown malignancy. This occurs in individuals who are immuno-suppressed intentionally, particularly children receiving organ transplants who are treated with cyclosporine A, or opportunistically, as in the case with individuals infected with HIV, or genetically, as in the case of affected males carrying the XLP (x-linked proliferative syndrome) gene. In these cases the resulting malignancies derive from the polyclonal proliferation of EBV-infected B cells. In addition, in such patients uncontrolled epithelial replication of the virus is detectable in lesions of oral hairy leukoplakia. Thus, the immune response plays a central role in the control of EBV infection.

As mentioned above EBV is a member of the herpesviruses. It possesses the following structural properties:

- The EBV genome consists of a linear double stranded DNA molecule (173,000 basepairs).
- The virion consists of a core (proteins and DNA), surrounded by an icosahedral capsid, and a membrane envelope enclosing the capsid. The icosahedral capsid is built up of hexameric and pentameric capsomeres. The membrane envelope consists of a protein/lipid bilayer membrane with spikes on its outer surface. The space between the capsid shell and the envelope is filled with amorphous protein, called the tegument.
- Like all herpesviruses, EBV is capable of establishing a latent life-long infection in its host subsequent to primary infection. This latency represents a perfect balance between EBV and its human host, controlled by the hosts immune system.

To date most biochemical and biological studies have been performed on three prototype strains of EBV, being B95-8 (transforming virus produced in a marmoset cell line), P3HR1 (non-transforming virus produced by a Burkitt's lymphoma tumor cell line) and Raji (latent virus in a Burkitt's lymphoma tumor cell line).

During the last few years the entire DNA sequence of prototype virus strain, B95-8, has been determined.

Analysis of this sequence has resulted in the identification of more than 80 open reading frames (Baer et al., 1984, Nature 310, p. 207-211).

The biology of EBV poses a special problem to investigators because its biological characteristics (latent infection) do not lend itself to the classic virus analysis. Furthermore, its cell and host range are effectively limited to human (and those of a few higher primates) B-lymphocytes and epithelial cells which are generally not amenable to culture in vitro. In addition, the absence of a fully permissive cell type, one in which the virus lytically replicates, has severely limited the ability to produce large amounts of the virus.

DNA molecules of B95-8, P3HR1- and Raji-isolates have been the prototypes for detailed restriction endonuclease mapping, and for cloning into Escherichia coli (E.coli) plasmids and in bacteriophage lambda, and for nucleotide sequencing.

The EBV-genome consists of a single double stranded DNA molecule build-up with unique and tandemly repeated DNA-elements. Each end of the DNA molecule contains multiple terminal sequences which permit covalently linking and circularization of the genome. In virus particles the EBV-genome is only detectable in a linear form. On the contrary, it exist as a circular episome inside the nucleus of latently infected cells.

The internal repeat sequences, IR1 to IR4, separate the EBV-genome into 5 unique regions. The U2 and U3 regions vary extensively among different EBV isolates and, the former being almost entirely deleted in the P3HR-1 strain of EBV.

The nomenclature for EBV reading frames is based on their position in the virus genome. The names begins with the initials of the BamH1 or EcoR1 restriction fragment where expression begins. The third letter in the name is L or R, depending on whether the expression is leftward or rightward on the standard map. (So BLLF2 is the second leftward reading frame starting in BamH1 restriction fragment L.).

The serological classification of virus antigens in the productive cycle of EBV is based on different fluorescence techniques.

Antigens specifically detected by means of the anti-complement immunofluorescence technique in the nucleus of fixed latently infected B-cells (e.g. Raji-cells) are classified as Epstein-Barr nuclear antigens (EBNA).

Upon activation of viral gene expression by chemical or viral factors a class of early antigens (EA) is detected whose synthesis is not blocked by inhibition of viral DNA synthesis. Dependent on the type of fixative used (Methanol or Acetone) two distinct sets of EA are detectable, EA_R and EA_D. EA is detectable by indirect immunofluorescence in the cytoplasm and nucleus of induced cells. Following onset of viral DNA-synthesis (and depending upon it) virus structural proteins (VCA) are synthesized which are detectable by indirect immunofluorescence in the cytoplasm and nucleus of virus producer cells (e.g. P₃HR₁ cells). On the surface of viable infected cells, induced for virus production a set of antigens (MA) is detectable by indirect immunofluorescence. These antigens can also be found on the viral envelope and are important targets for virus neutralization. Detection of EBV-specific antibodies in human sera can routinely be performed by serological techniques as described by Menke and Henle (Human Pathology, 5, 551-565, 1974).

Based upon biochemical and immunofluorescence data it is possible to distinguish five different classes of antigen molecules. The different viral polypeptides are designated by their molecular weight, and no common nomenclature has been established except for the virus envelope proteins.

The five different groups of antigens are:

- A. The group of antigens which are expressed during a state of latency (EBNAs and LMPs).
- B. The group of antigens which are responsible for genome activation and initial induction of viral replication (IEA).
- C. The group of antigens which are induced by IEA-gene products and which are required for replication of viral DNA; these antigens are mostly viral enzymes (EA).
- D. The group of antigens which are structural components of the viral particle and are expressed late in the viral replication cycle (VCA), after initiation of viral DNA-synthesis.
- E. The group of antigens which are expressed in the cell membrane of the infected cell (MA).

The viral capsid antigens (VCA) of EBV

For this antigen complex it also concerns that comparison of EBV specific proteins identified in different studies is difficult because of variations in polyacrylamide gelsystems, cell lines and chemical inducers used and the sera employed.

Dolyniuk et al. (1979) described a total of 33 proteins associated with purified virions. Differential solubilization with detergents suggest that the nucleocapsid is composed of at least seven proteins. An important component of the VCA complex is the major capsid protein (MCP). The EBV-MCP is encoded within the BcLF1 reading frame of the viral genome (Bear et al., 1984) and expressed as a 153-160 kDa non-glycosylated protein in EBV-producer cell lines with a pI of 7.5 to 9.0. This protein is synthesized in the cytoplasm in a soluble form and then transported to the nucleus, where it condenses into capsids and is no longer solubilized by detergents.

Another major VCA component has a molecular weight of 125 kDa and is glycosylated. This protein is encoded within the BALF4 reading frame of the viral genome. Although this glycoprotein was classified originally as a VCA component recent findings indicate that it might in fact be associated with cytoplasmic and nuclear membrane structures.

Experiments described previously (J.M. Middeldorp and P. Herbrink, J. Virol. Meth., 21, 133-146, 1988) aimed at the identification and characterization of diagnostically relevant EBV marker proteins in relation to different EBV-diseases.

This was done by using immunoblot strips containing antigens prepared from the virus producer cell line HH514-C16 (a superinducible derivative of P3HR1), induced for the expression of VCA/EA or EA, and from the EBV negative cell lines Ramos and Bjab. Cell lines which carry the EBV genome in a (fully) latent state,

X50-7 and JC-5, can be used to study EBNA/LMP specifically.

Patterns of EBV antibody responses were studied in sera of healthy seropositive blood donors, in sera of IM patients and chronic IM patients or patients with EBV-associated tumours like nasopharyngeal carcinoma. Polyclonal and monoclonal antibodies reactive with defined EBV-genome products can be used to characterize some of the protein bands detected in this experimental system. These studies however only described proteins or polypeptides with a certain molecular weight. No information was available as to the coding sequence on the EBV genome for these proteins. Nor was it known whether immunoreactive bands on immunoblots were due to reactivity with single or multiple proteins of the same molecular weight.

With immunoblot technique it is possible to detect an EBV antigen with a molecular weight of 18 kDa. This protein is not expressed when phosphono acetic acid (PAA) is used to block viral DNA-synthesis and is detected by all sera which contain anti-VCA antibodies which indicates that it is a VCA-related component. Another VCA component is a protein with a molecular weight of 40 kDa. Many of the viral capsid antigens are associated with the nuclear pellet.

At present EBV specific serodiagnosis is accomplished by rather subjective immunofluorescence tests. Progress to more simple and uniform diagnosis (e.g. ELISA) is hampered because bulk production and purification of viral antigens are not possible using standard virus producing cell lines. The only way to achieve this would be to use alternatively prepared EBV antigen(s). These EBV antigens could be prepared with either genetic engineering techniques or synthetic peptide techniques.

For the development of a specific and sensitive method to enable a reliable diagnosis to be made in various phases of the infection with EBV it is of great importance to identify immuno-dominant viral proteins and epitopes thereof.

The present invention provides peptides comprising at least part of the VCA-p18 or VCA-p40 protein, encoded within the EBV open reading frames BFRF3 and BdRF1 respectively, and fragments thereof, immunochemically reactive with antibodies to the Epstein Barr Virus. Part of the invention are therefore peptides with 176 and 345 amino acids respectively and an amino acid sequence as shown in SEQ ID NO: 2 and 4 which are immunochemically reactive with EBV antibodies.

The peptides according to the invention are found to be particularly suitable for use in a diagnostic method for the determination of the presence of EBV or EBV-antibodies in a sample. Moreover, a peptide according to the invention may be used in suitable pharmaceutical dosage forms in the treatment of an EBV-related disease. The preparation of vaccines thus obtained which contain a peptide or fragment thereof as active ingredients, is known to one skilled in the art.

In contrast to the natural EBV, the peptides according to the invention have the great advantage that these are of a safe non-infectious origin.

The invention also comprises fragments of said peptides which are still immunochemically reactive with antibodies to the Epstein-Barr Virus.

The term "peptide" as used herein refers to a molecular chain of amino acids with a biological activity, and does not refer to a specific length of the product. Thus inter alia, proteins, fusion-proteins or -peptides oligopeptides and polypeptides are included. If required peptides according to the invention can be modified in vivo or in vitro, for example by glycosylation, amidation, carboxylation or phosphorylation. Functional variants like, for example, acid addition salts, amides, esters, and specifically C-terminal esters, and N-acyl derivatives of the peptides according to the invention are therefore also considered part of the present invention. It will be understood that for the particular proteins or polypeptides embraced herein, natural variations can also exist. These variations may be demonstrated by (an) amino acid difference(s) in the overall sequence or by deletions, substitutions, insertions, inversions or additions of (an) amino acid(s) in said sequence. Amino acid substitutions from which can be expected that they do not essentially alter biological and immunological activities, have been described. Amino acid replacements between related amino acids or replacements which have occurred frequently in evolution are, inter alia Ser/Ala, Ser/Gly, Asp/Gly, Asp/Asn, Ile/Val (see Dayhof, M.D., Atlas of protein sequence and structure, Nat. Biomed. Res. Found., Washington D.C., 1978, vol. 5, suppl. 3). Based on this information Lipman and Pearson developed a method for rapid and sensitive protein comparison (Science 227, 1435-1441, 1985) and determining the functional similarity between homologous proteins.

The term "fragment" as used herein means an amino acid sequence comprising a subsequence of a peptide of the invention. Said fragment is a peptide having one or more immunogenic determinants of the VCA-p18 or VCA-p40 protein. Fragments can inter alia be produced by enzymatic cleavage of precursor molecules, using restriction endonucleases for the DNA and proteases for the polypeptides. Other methods include chemical synthesis of the fragments or the expression of peptide fragments by DNA fragments.

Suitable immunogenic fragments of a peptide according to the invention containing (an) epitope(s) can be found by means of the method described in Patent Application WO 86/06487, Geysen, H.M. et al. (Proc.

Natl. Acad. Sci. 81, 3998-4002, 1984), Geysen, H.M. et al. (J. Immunol. Meth. 102, 259-274, 1987) based on the so-called pepscan method, wherein a series of partially overlapping peptides corresponding with partial sequences of the complete polypeptide under consideration, are synthesized and their reactivity with antibodies is investigated.

5 In addition, a number of regions of the peptides can be designated epitopes on the basis of theoretical considerations, although the predictive value of these theoretical considerations is limited. The determination of these regions is based on a combination of the hydrophilicity criteria according to Hopp and Woods (Proc. Natl. Acad. Sci. 78, 3824-3828, 1981) and the secondary structure aspects according to Chou and Fasman (Advances in Enzymology 47, 45-148, 1987).

10 Preferred peptides according to the invention are peptides comprising at least one of the amino acid sequences as shown in SEQ ID NO:5 and SEQ ID NO:6.

Most preferred is a peptide comprising the amino acid as shown in SEQ ID NO:5 linked to the amino acid sequence as shown in SEQ ID NO:6. Such a combi-peptide has proven to be extremely useful for the specific detection of IgG, IgA, IgM antibodies to EBV-VCA, with sensitivity similar or even better than standard serological techniques. As such IgM-EBV is a useful diagnostic marker for acute primary EBV-infection, whereas IgA to EBV is useful for diagnosis and prognosis in Nasopharyngeal Carcinoma. EBV-IgG is positive in all human EBV-carriers and negative in persons not infected with the virus. In addition changes in the antibody titer for each of the antibodies of a specific subclass may be of additional diagnostic value. Since antibodies of different subclasses have a specific diagnostic value in different stages of EBV-infection 15 the use of a combi-peptide according to the invention in diagnostic tests, e.g. ELISA, can be of great advantage.

The preparation of the peptides or fragments thereof according to the invention is effected by means of one of the known organic chemical methods for peptide synthesis or with the aid of recombinant DNA techniques.

25 The organic chemical methods for peptide synthesis are considered to include the coupling of the required amino acids by means of a condensation reaction, either in homogeneous phase or with the aid of a so-called solid phase.

The condensation reaction can be carried out as follows:

30 a) condensation of a compound (amino acid, peptide) with a free carboxyl group and protected other reactive groups with a compound (amino acid, peptide) with a free amino group and protected other reactive groups, in the presence of a condensation agent;

b) condensation of a compound (amino acid, peptide) with an activated carboxyl group and free or protected other reaction groups with a compound (amino acid, peptide) with a free amino group and free or protected other reactive groups.

35 Activation of the carboxyl group can take place, inter alia, by converting the carboxyl group to an acid halide, azide, anhydride, imidazolide or an activated ester, such as the N-hydroxy-succinimide, N-hydroxy-benzotriazole or p-nitrophenyl ester.

The most common methods for the above condensation reactions are: the carbodiimide method, the azide method, the mixed anhydride method and the method using activated esters, such as described in 40 The Peptides, Analysis, Synthesis, Biology Vol. 1-3 (Ed. Gross, E. and Meienhofer, J.) 1979, 1980, 1981 (Academic Press, Inc.).

Preparation of suitable fragments of above-mentioned peptides according to the invention using the "solid phase" is for instance described in J. Amer. Chem. Soc. 85, 2149 (1963) and Int. J. Peptide Protein Res. 35, 161-214 (1990). The coupling of the amino acids of the peptide to be prepared usually starts from 45 the carboxyl end side.

For this method a solid phase is needed on which there are reactive groups or on which such groups can be introduced. This can be, for example, a copolymer of benzene and divinylbenzene with reactive chloromethyl groups, or a polymeric solid phase rendered reactive with hydroxymethyl or amine-function.

A particularly suitable solid phase is, for example, the p-alkoxybenzyl alcohol resin (4-hydroxy-methyl-phenoxy-methyl-copolystyrene-1% divinylbenzene resin), described by Wang (1974) J. Am. Chem. Soc. 95, 1328. After synthesis the peptides can be split from this solid phase under mild conditions.

After synthesis of the desired amino acid sequence, detaching of the peptide from the resin follows, for example, with trifluoromethanesulphonic acid or with methanesulphonic acid dissolved in trifluoroacetic acid. The peptide can also be removed from the carrier by transesterification with a lower alcohol, preferably 55 methanol or ethanol, in which case a lower alkyl ester of the peptide is formed directly. Likewise, splitting with the aid of ammonia gives the amide of a peptide according to the invention.

The reactive groups which may not participate in the condensation reaction are, as stated, effectively protected by groups which can be removed again very easily by hydrolysis with the aid of acid, base or

reduction. Thus, a carbonyl group can be effectively protected by, for example, esterification with methanol, ethanol, tertiary butanol, benzyl alcohol or p-nitrobenzyl alcohol and amines linked to solid support.

Groups which can effectively protect an amino group are the ethoxycarbonyl, benzyloxycarbonyl, t-butoxy-carbonyl (t-boc) or p-methoxy-benzyloxycarbonyl group, or an acid group derived from a sulphonic acid, such as the benzene-sulphonyl or p-toluene-sulphonyl group, but other groups can also be used, such as substituted or unsubstituted aryl or alkyl groups, for example benzyl and triphenylmethyl, or groups such as ortho-nitrophenyl-sulphenyl and 2-benzoyl-1-methyl-vinyl. A particularly suitable α -amino-protective group is, for example, the base-sensitive 9-fluorenyl-methoxycarbonyl (Fmoc) group [Carpino & Han (1970) J. Amer. Chem. Soc. 92, 5748].

A more extensive account of possible protecting groups can be found in *The Peptides, Analysis, Synthesis, Biology*, Vol. 1 - 9 (Eds. Gross, Udénfriend and Meienhofer) 1979 - 1987 (Academic Press, Inc.).

It is necessary also to protect the ϵ -amino group of lysine and advisable for the guanidine group of arginine. Customary protective groups in this connection are a Boc-group for lysine and a Pmc- or Pms- or Mbs-group or Mtr-group for arginine.

The protective groups can be split off by various conventional methods, depending on the nature of the particular group, for example with the aid of trifluoroacetic acid or by mild reduction, for example with hydrogen and a catalyst, such as palladium, or with HBr in glacial acetic acid.

As already indicated above, the peptides according to the invention can likewise be prepared with the aid of recombinant DNA techniques. This possibility is of importance particularly when the peptide is incorporated in a repeating sequence ("in tandem") or when the peptide can be prepared as a constituent of a (much larger) protein or polypeptide or as a fusion protein with, for example, (part of) β -galactosidase. This type of peptides therefore likewise falls within the scope of the invention. For this purpose, as a constituent of a recombinant DNA, a nucleic acid sequence is used which codes for a peptide according to the invention and which, furthermore, is substantially free from nucleic acid segments, which in the naturally occurring EBV genome flank the nucleic acid sequence indicated above.

This latter method involves the preparation of the desired peptide by means of bringing to expression a recombinant polynucleotide with a nucleic acid sequence which is coding for one or more of the peptides in question in a suitable micro-organism as host.

The invention therefore further encompasses nucleic acid sequences encoding a peptide according to the invention, preferably comprising at least part of the nucleic acid sequence as shown in SEQ ID NO: 1 and/or 3.

"Nucleic acid sequence" as used herein refers to a polymeric form of nucleotides of any length, both to ribonucleic acid sequences and to deoxy ribonucleic acid sequences. In principle, this term refers to the primary structure of the molecule. Thus, this term includes double and single stranded DNA, as well as double and single stranded RNA, and modifications thereof.

A nucleic acid sequence according to the present invention can be ligated to various replication effecting DNA sequences with which it is not associated or linked in nature resulting in a so called recombinant vector molecule which can be used for the transformation or transfection of a suitable host. Useful recombinant vector molecules, are preferably derived from, for example plasmids, bacteriophages, cosmids or viruses.

Specific vectors or cloning vehicles which can be used to clone nucleic acid sequences according to the invention are known in the art and include inter alia plasmid vectors such as pBR322, the various pUC, pGEM and Bluescript plasmids, bacteriophages, e.g. λ gt-Wes, Charon 28 and the M13 derived phages or viral vectors such as SV40, adenovirus or polyoma virus (see also Rodriguez, R.L. and D.T. Denhardt, ed., Vectors: A survey of molecular cloning vectors and their uses, Butterworths, 1988; Lenstra, J.A. et al., Arch. Virol. 110, 1-24, 1990). The methods to be used for the construction of a recombinant vector molecule according to the invention are known to those of ordinary skill in the art and are inter alia set forth in Maniatis, T. et al. (Molecular Cloning A Laboratory Manual, second edition; Cold Spring Harbor Laboratory, 1989).

For example, the insertion of the nucleic acid sequence according to the invention into a cloning vector can easily be achieved when both the genes and the desired cloning vehicle have been cut with the same restriction enzyme(s) as complementary DNA termini are thereby produced.

The recombinant vector molecules according to the invention may additionally contain one or more marker activities that may be used to select for desired transformants, such as ampicillin and tetracycline resistance in pBR322, as for example ampicillin resistance and α -peptide of β -galactosidase in pUC8.

The invention also comprises (a) host cell(s) transformed or transfected with a nucleic acid sequence or recombinant expression vector molecule described above, capable of producing the peptides according to the invention by expression of the corresponding nucleic acid sequence.

A suitable host cell is a microorganism or cell which can be transformed by a nucleic acid sequence encoding a peptide or by a recombinant vector molecule comprising such a nucleic acid sequence and which can if desired be used to express said peptide encoded by said nucleic acid sequence. The host cell can be of procaryotic origin, e.g. bacteria such as *Escherichia coli*, *Bacillus subtilis* and *Pseudomonas* species; or of eucaryotic origin such as yeasts, e.g. *Saccharomyces cerevisiae* or higher eucaryotic cells such as insect, plant or mammalian cells, including HeLa cells and Chinese hamster ovary (CHO) cells. In general, prokaryotes are preferred for the construction of the recombinant vector molecules useful in the invention. For expression nucleic acid sequences of the present invention are introduced into an expression vector, i.e. said sequences are operably linked to expression control sequences. Such control sequences may comprise promoters, enhancers, operators, inducers, ribosome binding sites etc. Therefore, the present invention provides a recombinant vector molecule comprising a nucleic acid sequence encoding the peptides identified above operably linked to expression control sequences, capable of expressing the DNA sequences contained therein in (a) transformed or transfected host cell(s).

It should, of course, be understood that the nucleotide sequences inserted at the selected site of the cloning vector may include only a fragment of the complete nucleic acid sequence encoding for the peptides according to the invention as long as the transformed or transfected host will produce a polypeptide having at least one or more immunogenic determinants.

Antibodies, directed to a peptide according to the invention are also part of the present invention.

The peptides or fragments thereof prepared and described above are used to produce antibodies, both polyclonal and monoclonal. Monoclonal antibodies directed against peptides according to the invention can be readily produced by one skilled in the art.

Preferred antibodies to different epitopes of the VCA-p18 protein according to the invention are antibodies having the same reactivity with VCA-p18 as antibodies produced by the rat-mouse hybridoma cell line deposited at the European Culture of Animal Cell Cultures (ECACC), Porton Down (UK), under the deposit no.s 93020413 or 93020412. Preferred antibodies to an epitope of the VCA-p40 protein are antibodies having the same reactivity with VCA-p40 as antibodies produced by the mouse-mouse hybridoma cell line deposited at the European Culture of Animal Cell Cultures (ECACC), Porton Down (UK), under the provisional deposit no. 93020414.

Immortalized cell lines capable of excreting monoclonal antibodies according to the invention are also part of the present invention.

The preparation of cell lines producing monoclonal antibodies may occur by, for example, by the Kohler and Milstein technique (Kohler and Milstein devised the techniques that resulted in the formation monoclonal antibody-producing hybridomas (G. Kohler and C. Milstein, 1975, *Nature* 256:495-497; 1976, *Eur. J. Immunol.* 6:511-519)), transformation with Epstein-Barr Virus, or a direct transformation technique of B-lymphocytes with oncogenic DNA, or a direct fusion of human B-lymphocytes with a fusion partner being either a human or a mouse-human hybrid myeloma cell line, or a direct fusion of an EBV-transformed B cell line with said myeloma cell lines.

Preferred cell lines according to the invention are the cell lines deposited at the European Collection of Animal Cell Cultures, Porton Down (UK) under deposit No.s 93020413, 93020412, 93020414. These cell lines have been deposited at the ECACC on 4 February, 1993, under the terms and conditions of the Budapest treaty, 1977.

The cell lines with deposit no.s. 93020413 and 93020412 are both capable of producing monoclonal antibodies (EBV.OT15E and EBV.OT15I respectively) to two respective epitopes of the VCA-p18 protein. These cell lines are rat-mouse hybridoma cell lines. The cell line with provisional deposit no. 93020414 is capable of producing antibodies to an epitope of the VCA-p40 protein, and is a mouse-mouse hybridoma cell line. The antibodies produced by these cell lines have been used to identify epitopes on the respective proteins (as is further exemplified in the examples).

Antibodies, both monoclonal and polyclonal, directed against peptides according to the invention are very suitable in diagnosis and immunocytochemistry for detection in situ in tissue specimen, while those antibodies which are neutralizing are very useful in passive immunotherapy. Especially monoclonal antibodies may be used to raise anti-idiotypic antibodies. Techniques for raising anti-idiotypic antibodies are known in the art. Anti-idiotypic antibodies reactive with the monoclonal antibodies according to the invention, as described above, are part of the present invention.

Anti-idiotypic antibodies are antibodies directed to the variable part of immunoglobulins. A sub-population of anti-idiotypic antibodies is known as "anti-idiotypic β " or "internal images". These anti-idiotypic β antibodies have either a structural or a three dimensional resemblance with the antigen (Uytendaele, F.G.C.M. et al. *Immunol.Rev.* 90: 93-113; 1986). This type of anti-idiotypic antibodies is widely used as a vaccine against infectious diseases in animal models (Hiernaux J.R.; *Infect.Immun.*; 56; 1407-1413; 1988,

Kennedy, R.C. et al.; Science 232: 220-223;1986). For use in assays the anti-idiotypic antibodies can be raised in large amounts.

Techniques for raising anti-idiotypic antibodies are known in the art. For example, anti-idiotypic antibodies according to the invention can be obtained by immunizing BALB/c mice with monoclonal antibodies, coupled to KLH with glutaraldehyde according to standard literature procedures, mixed with Freund's complete adjuvant. The spleen cells of these mice can be immortalized and the thus obtained hybridomas can be screened for anti-idiotypic antibody production. Screening of the hybridomas can be performed, for example, by binding monoclonal antibodies according to the invention to a solid phase (wells of microtiter plates) and incubating the solid phase with culture supernatant of growing hybridomas. An EBV peptide coupled to Horse Radish Peroxidase (HRP) can be added. The presence of anti-idiotypic antibodies in culture supernatant will then be indicated by inhibition of the binding of this peptide conjugate to the monoclonal antibodies coated on the solid phase.

Anti-idiotypic antibodies can be used for instance for inhibiting the binding of human and/or animal EBV-antigen in an immuno assay using EBV-antibodies. Alternatively anti-idiotypic antibodies can be used as a mimicing agent of the immunochemical reagent mentioned hereunder.

Said anti-idiotypic antibodies are also useful for diagnosis and treatment of EBV, as well as for the elucidation of important epitopic regions of EBV-antigens.

An immunochemical reagent comprising one or more peptides or antibodies according to the invention is also part of the present invention.

The term "immunochemical reagent" according to the invention usually consists of one more peptides according to the invention and a suitable support or a labelling substance.

Supports which can be used are, for example, the inner wall of a microtest well or a cuvette, a tube or capillary, a membrane, filter, test strip or the surface of a particle such as, for example, a latex particle, an erythrocyte, a dye sol, a metal sol or metal compound as sol particle, a carrier protein such as BSA or KLH. Labelling substances which can be used are, inter alia, a radioactive isotope, a fluorescent compound, an enzyme, a dye sol, metal sol or metal compound as sol particle.

In a method for the detection of antibodies directed against EBV in a sample, an immuno-chemical reagent according to the invention is brought into contact with the sample. After which, the presence of immune complexes formed between the peptide and antibodies in the sample is detected and by this detection the presence of EBV antibodies in the sample is known and can be determined quantitatively.

Depending on the nature and further characteristics of the immunochemical reagent the immunochemical reaction that takes place is a so called sandwich reaction, an agglutination reaction, a competition reaction or an inhibition reaction.

For the detection of EBV in a sample an immunochemical reagent according to the invention, containing one or more peptides according to the invention, can be brought into contact with the sample and anti-EBV after which the presence of immune complexes formed can be detected and, from this, the presence of EBV in a sample can be determined.

A particularly suitable method for the detection of EBV in a sample is based on a competition reaction between a peptide according to the invention provided with a labelling substance and an EBV antigen (present in the sample) whereby the peptide and the antigen are competing with the antibody directed against EBV attached to a solid support.

The invention further comprises a method for the detection of Epstein-Barr virus in a sample characterized in that an antibody according to the invention is brought into contact with a sample whereafter the presence of immune complexes formed is detected which is a measure for the presence of Epstein barr Virus in the sample.

A test kit according to the invention comprises as an essential constituent an immunochemical reagent as described above. Carrying out a sandwich reaction, for the detection of EBV antibodies the test kit may comprise, for example, the peptide according to the invention coated to a solid support, for example the inner wall of a microtest well, and either a labelled peptide according to the invention or a labelled anti-antibody.

For carrying out a competition reaction, the test kit may comprise a peptide according to the invention coated to a solid support, and a labelled antibody directed against EBV preferably a monoclonal antibody directed against said peptide.

In an agglutination reaction the test kit comprises an immunochemical reagent which may comprise a peptide according to the invention coated to particles or sols. Another embodiment of a test kit is, for example, the use of a labelled peptide according to the invention as immunochemical reagent in a competition reaction with an EBV antigen to be detected for a binding site on the antibody directed against EBV, which is coated to a solid support.

It is within the scope of this invention to use the new nucleotide sequences according to SEQ ID NO: 1 and/or 3 as the basis of a test to detect EBV DNA or RNA by a nucleic acid amplification technique for instance the polymerase chain reaction (PCR) or the nucleic acid sequence based amplification (NASBA), as described in EP 201,814 and EP 329,822, respectively. A method for the amplification and the detection of an Epstein-Barr Virus nucleic acid sequence in a sample using at least one nucleic acid sequence or fragment thereof according to the invention primer(s) in order to perform a nucleic acid amplification of said Epstein-Barr Virus nucleic acid sequence and to detect the amplified sequence is also part of the present invention. Part of the invention is also a test amplification kit for carrying out above-mentioned amplification technique, said kit containing at least a set of primers corresponding to at least a part of the nucleotide sequences according to the invention.

The invention is further exemplified by the following examples:

LEGENDS:

Figure 1: Outline of the approach used to identify the EBV-encoded genes coding for the VCA-p40 and VCA-p18 proteins.

Figure 2:

- a). Western blot of nuclear antigen extract of virus-producer cell line HH514, induced for expression of EA and VCA.
 - b) Western blot of whole cell lysate of E.coli expressing the BFRF3- β -galactosidase fusion protein.
 - c) Western blot of whole cell lysate of E.coli expressing the BDRF1- β -galactosidase fusion protein.
 - d) Western blot of whole cell lysate of E.coli expressing β -galactosidase only.
- The blots were probed with a set of individual sera. Sera used in lanes 1-16 of the blots were respectively:
1. mouse monoclonal antibody to β -Gal (Promega)
 2. mouse monoclonal antibody to VCA-P40 raised by immunizing with natural viral capsid proteins (EBV.OT41A)
 3. Human antibody mono-specific for viral VCA-P18 obtained by specific immuno affinity purification with viral VCA-P18
 - 4-5 Human EBV-seronegative sera
 - 6-16 Human EBV-seropositive sera with a different relative reactivity towards viral VCA-P18 and VCA-P40

Figure 3:

- Immunoblots of three human sera (serum 92, serum 214 and serum 219) on nitrocellulose strips pre-absorbed with the following amounts of BFRF3- β -galactosidase fusion protein:
- lane 1: 0 μ g BFRF3- β -galactosidase
 - lane 2: 0.01 μ g BFRF3- β -galactosidase
 - lane 3: 0.1 μ g BFRF3- β -galactosidase
 - lane 4: 0.5 μ g BFRF3- β -galactosidase
 - lane 5: 1 μ g BFRF3- β -galactosidase
- The lane in which β -galactosidase only is present is indicated as lane β .

Figure 4:

Results of PEPSCAN analysis: Percentage of 15 human sera from healthy EBV-seropositive donors reactive with 12-mer peptides from the VCA-18 sequence. Starting positions of the 12-mer peptides within the amino acid sequence of the VCA-p18 protein are indicated on the X-axis.

Figures 5a and 5b:

PEPSCAN results (optical density at 450 nm) of an analysis of VCA-p18 derived 12-mer peptides, using two rat monoclonal antibodies directed against VCA-p18 (EBV.OT15E and EBV.OT15I respectively).

Figure 6:

PEPSCAN results of an analysis of 12-mer peptides derived from the VCA-p40 protein with a mouse monoclonal antibody (EBV.OT41A) directed against VCA-p40.

Figure 7:

ELISA reactivity (optical density at 450 nm) of 43 human serum samples obtained from healthy blood donors tested for IgG-reactivity against selected synthetic peptides derived from the BFRF3-encoded VCA-p18 protein.

- Δ indicates sera negative by standard serological analysis
- indicates sera positive by standard serological analysis
- ◇ indicates sera indeterminate for EBV antibodies by standard serological analysis but negative on immunoblot.
- * indicates sera positive by standard serology but negative for anti-p18 antibodies on immunoblot.

Peptide 1: H₂N-GVPRRQRAIDKRQRA-COOH

Peptide 2: H₂N-GQPHDTAPRGARKKQ-COOH

Peptide 3: H₂N-AVDTGSGGGGQPHDTAPRGARKKQ-COOH

Peptide 4: H₂N-STAVAQSATPSVSSSISSLRAATSGATAAA-COOH

Peptide 5: Combi-peptide of peptide 4 and 3 linked by S-S-bridging.

Figure 8 a, b, c:

Analysis of immunoreactivity (ELISA) of the combi peptide no.5 with:

- a) human IgG (76 VCA-I.F. positive sera)
- b) human IgM (26 IgM positive sera)
- c) human IgA (35 sera from NPC patients)

EXAMPLES**Example 1**

Procedure for isolating and identifying new DNA sequences coding for EBV proteins.

The general strategy followed for the identification resulting in the new EBV marker molecules can be subdivided in the following phases.

1. Identification and production of antibody reagents specifically reactive with the EBV marker molecules.
2. Preparing a c-DNA bank from poly-A selected or total cell mRNA, isolated from EBV expressing cells, or preparing a genomic DNA-bank from fragments of the viral genome, preferably in phage lambda gt11, followed by screening of the phage synthesized proteins with above antibody reagents.
3. Purification of reactive phages and identification of the EBV-specific insert sequences contained within the genome of the phages.
4. Correlation of the insert sequences with the published prototype EBV-genome sequence to locate the corresponding EBV-specified open reading frames.
5. Cloning, expression and production of identified open reading frames in alternative host cells such as E.coli, Baculovirus, yeast or higher eukaryotic cells or alternative expression c.q. production systems.

These procedures are outlined in detail below and illustrated by the scheme as depicted in figure 1.

(Phase 1:)Cell cultures and cell extracts.

The P3HR1-derived cell line HH514.c16 was propagated as a suspension culture in roller bottles and induced for VCA- and EA-expression using 20ng/ml 12-tetradecanophorbol-13-acetate (TPA) and 3mM sodium butyrate exactly as described by Middeldorp and Herbrink (J.Virol.Neth., 21, 133-146, 1988). For the selective expression of EA-antigens only, viral DNA polymerase was blocked by the addition of 0.5mM phosphonoacetic acid to the induced cell culture.

Monoclonal antibodies.

For the production of monoclonal antibodies BALB/c mice were immunized with the nuclear fraction of VCA-induced HH514 cells (F.Wielaard et al., J.Virol.Meth., 21, 105-115, 1988) or with more extensively purified proteins from these cells or from alternative expression systems. Hybridomas were produced according to standard protocols and supernatants were analysed in standard EBV immunofluorescence tests and on immunoblot strips containing antigen extracts from VCA-induced HH514 cells as described by Middeldorp and Herbrink, J.Virol.Meth., 21, 133-146, 1988).

Affinity purification of anti VCA-p18 antibodies.

Anti VCA-p18 antibodies were purified from a human EBV-positive serum according to the method of Robinson and Miller (The herpesviruses, Vol.1, 151-192, 1991, Publ.Plenum Publishing Corp. New York) with some minor modifications. In short, following the electrophoretical separation in 10% acrylamide gels, the proteins were blotted onto PVDF-membranes (Millipore Corporation, Bedford, USA), and the region on the PVDF-membrane corresponding to VCA-p18 was cut out and used as affinity matrix. Nonspecific binding of antibodies to the strips was prevented by incubation overnight in blocking solution (5% dried milk-powder, 4% horse serum in phosphate buffered saline, pH 7.4 (PBS)). Thereafter the strips were incubated with diluted human serum (1:25 in blocking solution) for 2 hours. After three wash steps with PBS containing 0.05% Tween-20 the bound antibodies were eluted with 0.1 M glycine, pH 2.7 in two consecutive incubations. The eluate was neutralized with 1/20th volume of 1M Tris/HCl, pH 9.0. Finally, the eluate was dialyzed against PBS and stored in aliquots at -20°C.

(Phase 2:)

RNA purification of HH514.c16 cells.

Total RNA was isolated from induced HH514.c16 cells by the guanidinium/CsCl procedure as described by Maniatis et al. (Molecular cloning: A laboratory manual; Cold Spring Harbor Laboratory, New York, 1982). Purification of poly (A⁺) RNA was performed by oligo(dT) chromatography (Pharmacia, Inc., Piscataway N.Y.) as described by Ausubel et al. (Current protocols in molecular biology (1991) Greene Publishing Associates, John Wiley & Sons, New York).

Northern blot analysis.

Total RNA (10 µg) was denatured by glyoxal (P.S.Thomas, Methods in Enzymology, 100, 255-266, 1983) and run in agarose gels. After ethidium bromide staining, the separated RNA was vacuum-blotted to nitrocellulose. After 3hr prehybridization the filter was hybridized overnight at 42 °C with [α -32P]-labeled (Amersham, Bristol, UK) randomly primed or nick translated DNA probes. The hybridization solutions consisted of 50% formamide, 5xSSPE, SxDenhardtts solution (0.1% (w/v) polyvinylpyrrolidone, 0.1% (w/v) BSA, 0.1% (w/v) Ficoll), 0.2 mg sonicated herring sperm DNA per ml and 0.5% SDS. Subsequently, blots were washed and exposed to X-ray film (Eastman Kodak Co., Rochester, N.Y., USA) using intensifying screens.

Constuction of cDNA library in lambda-gt11.

Five micrograms of poly (A⁺)-selected RNA was denatured with methylmercury hydroxide. The cDNA synthesis was primed with either oligo(dT) or hexanucleotides. Procedures for first and second-strand synthesis were identical to those described by Gubler and Hoffman (Gene, 25, 263-269, 1983). After EcoRI-methylation, EcoRI-linker addition and EcoRI digestion, the modified cDNA was size selected by sepharose CL4B (Pharmacia) chromatography as described in Maniatis et al. cDNA varying in size from 0.5kB to 3.0kB was ligated into phosphatase treated lambda-gt11-arms followed by in vitro packaging using the Packagene kit (Promega, San Diego, USA).

Immunological screening of the lambda-gt11 library.

A total of 1×10^4 recombinant phages of the oligo(dT) primed library and 5×10^5 recombinants of the hexanucleotide primed library were screened for immunological reactivity (see Maniatis et al.) with affinity

purified human anti-p18 antibodies or monoclonal antibody EBV.OT41A. Immunoreactive plaques were detected with alkaline phosphatase-conjugated anti-human or antimouse IgG, respectively, as described by the manufacturer (Promega).

5 (Phase 3:)

Nucleotide sequence analysis.

10 Insert DNA of positive plaques, purified by repeated plaque lifting and immunoscreening, was amplified by the polymerase chain reaction (PCR) technique using primers of the lambda-gt11 flanking sequences containing restriction sites at their 5'-end. After digestion with the appropriate restriction enzyme the DNA-fragment was subcloned in pGEM-4Z and sequenced from both sides using a sequence kit (Pharmacia, Uppsala, Sweden) which employs a modification on the method of Sanger et al. (Proc.Natl.Acad.Sci., USA, 74, 5463-5467, 1977).

15 (Phase 4:)

Alignment of sequences.

20 Sequences were aligned with the published sequence of EBV-B95-8 prototype (Baer et al., Nature 310, 207-211, 1984) as deposited at the EMBL-sequence database, using the software programs of the University of Wisconsin, genetics computer group (Gribshov et al. Nucl.Acid res., 14, 327-334, 1986).

25 (Phase 5:)

Cloning and expression of BFRF3 and BdRF1 in E.coli.

The EBV genome-encoded open reading frames (ORFs), BFRF3 and BdRF1, were amplified by PCR using viral DNA, purified from virions isolated by sucrose-density centrifugation from HH514.c16 culture supernatants as a target. Primers of each set containing restriction sites at their 5'-end were used for 30 cloning the amplified fragment in the EcoRI-HindIII site of expression vector pMLB1113 (a derivative of PBR322) (Zagurski and Berman, Gene, 27, 183-101, 1984) which is located at the fifth codon of the 5' end of the LacZ gene. Proteins expressed from these constructs consist of the first 5 amino acids of β -galactosidase followed by the recombinant protein linked at their C-terminus to the remainder of β -galactosidase. Non-fusion proteins were constructed similarly, but with a stopcodon inserted at the 3' end of the insert.

E.coli expression of recombinant proteins.

40 Transformed E.coli cultures were induced by addition of 1 mM IPTG (isopropyl- β -D-thiogalactopyranoside). At 2 hrs post-induction bacterial cells were collected by centrifugation and suspended in SDS-PAGE sample buffer and analysed by immunoblotting.

45 Example 2:

Immunoreactivity of the VCA-p18 and VCA-p40 proteins.

The immunoreactivity of the proteins obtained following the procedure as outlined in example 1 was investigated by immunoblot analysis. For this purpose whole cell proteins of either VCA-induced HH514.c16 50 cells or E.coli expressing BFRF3- or BdRF1- β -galactosidase fusion proteins or β -galactosidase only were separated by SDS-PAGE and blotted onto nitrocellulose. Strips prepared from these blots were incubated with individual sera and antibody preparations.

Immunoblot procedure:

55 The immunoblot procedure was performed essentially as described by Middeldorp and Herbrink (J.Virol.Meth., 21 p133-159, 1988) and is briefly outlined below: After SDS-PAGE proteins were transferred to nitrocellulose filters (0.2 μ Schleicher & Schuell, Den Bosch, the Netherlands). Nonspecific binding of

antibodies to the filters was prevented by incubation for at least 1 hr at room temperature (RT) with blocking buffer (4% dried milk powder, 5% horse serum in Tris buffered saline (TBS)). Human sera were diluted to an appropriate dilution in blocking buffer and incubated for at least 1 hr. Blots or strips were washed three times in TBS + 0.05% Tween-20 (TBSt), and alkaline phosphatase (AP)-conjugated anti-human IgG antibody (Promega), or HRP-conjugated anti-mouse IgG, or anti-rat IgG (Organon Teknika Cappel, Boxtel, the Netherlands) was added at the appropriate dilution in blocking buffer. After further incubation and rinse steps, the blot was developed using nitroblue tetrazolium (NBT) and 5-bromo-4-chloro-3 indolyl phosphate (BCIP) as substrates for AP and 4-chloro-naphtol for HRP.

The resulting blots are depicted in figure 2a-d. Sera used in lanes 1-12 of the blots were respectively:

1. mouse monoclonal antibody to β -Gal (Promega)
2. mouse monoclonal antibody to VCA-P40 raised by immunizing with natural viral capsid proteins (EBV.OT41A)
3. Human antibody mono-specific for viral VCA-P18 obtained by specific immuno affinity purification with viral VCA-P18
- 4-5 Human EBV-seronegative sera
- 6-16 Human EBV-seropositive sera with a different relative reactivity towards viral VCA-P18 and VCA-P40.

Figure 2a shows immunoreactivity of the natural viral polypeptides, separated by reducing SDS-PAGE in 12.5% acrylamide. Anti β -galactosidase antibodies are negative (1), whereas anti-sera specific for VCA-p40 (2) and VCA-p18 (3) react with the respective viral proteins. Strips 4 and 5, stained with human EBV-negative sera, show no reactivity, whereas strips 6-16 represent different immunoreactivities with EBV-proteins as found in normal healthy seropositive blood donors (6-15) or patients with active EBV disease (16). Positions of the VCA-p18 and VCA-p40 bands are indicated on the right hand side.

Figure 2b shows immunoreactivity of whole cell proteins from E.coli expressing the VCA-p18 β -galactosidase fusion protein (BFRF3-LacZ) separated by reducing SDS-PAGE in 10% acrylamide. Anti β -galactosidase (1) and anti VCA-p18 (3) antibodies specifically react with a number of protein bands which are due to proteolytic fragmentation in E.coli of the original fusion protein at 134 kDa. The anti-VCA-p40 monoclonal antibody (2) is negative, just like human EBV-negative sera (4-5). Human sera with different reactivities to viral VCA-p18 show similar different reactivities on the fusion protein. Again staining of multiple bands are due to proteolytic breakdown of the fusion protein in E.coli.

Figure 2c shows the same analysis for the VCA-p40- β -galactosidase fusion protein in E.coli, which is less sensitive to proteolysis resulting in a single band at 156kDa. Interpretation of figure 2c is analogous to 2b.

Figure 2d represents a control blot of E.coli expressing β -galactosidase only.

From the above described study it is apparent that the individual E.coli constructs indeed express the respective fusion proteins. Human sera show the same reactivity towards the expressed proteins as towards the natural viral proteins.

Example 3:

Immunoabsorption.

Immunochemical identity of an alternative peptide according to the invention and the corresponding natural viral protein can be proven by pre-absorption of human sera, reactive on immunoblots containing natural viral protein(s), with different concentrations of the peptide according to the invention, resulting in the disappearance of specific antibody reactivity to the corresponding natural viral protein on the blot. In addition to proving immunochemical identity, this technique provides strong evidence that immunoreactivity of a viral protein on immunoblots is mediated by antibody binding to a single (dominant) protein species.

An experiment as described above was performed for the VCA-p18 marker protein:

The immunoreactivity of human sera, (serum 92, serum 214 and serum 219), obtained from three healthy EBV-seropositive donors, was studied by immunoblot analysis on nitrocellulose strips containing nuclear antigens from VCA-induced HH514 cells, separated by reducing SDS-PAGE in 10% acrylamide gels. Sera were preabsorbed overnight at +4°C with increasing amounts of purified BFRF3- β -galactosidase fusion protein, or with β -galactosidase only before immunoblot analysis. (Staining for IgG-reactivity was performed with peroxidase-labeled sheep anti-human IgG), using N-chloro-naphtol as precipitating substrate.

The results of this experiment are illustrated by figure 3. The amounts of purified BFRF3- β -galactosidase fusion protein used in pre-absorption of 1 ml 1:100 diluted serum analysed in the respective lanes of the immunoblots as depicted in figure 3 were:

lane 1: 0 μ g BFRF3- β -galactosidase

- lane 2: 0.01 μ g BFRF3- β -galactosidase
 lane 3: 0.1 μ g BFRF3- β -galactosidase
 lane 4: 0.5 μ g BFRF3- β -galactosidase
 lane 5: 1 μ g BFRF3- β -galactosidase

5 The lane in which only β -galactosidase was used for preabsorption is indicated as lane β .

From figure 3 it can be seen that reactivity of the band representing VCA-p18 (arrow) disappears with increasing concentrations of BFRF3- β -galactosidase fusion protein during preabsorption. Preabsorption with β -galactosidase alone had no effect, whereas reactivity with other EBV-proteins is non-effected by any of the preabsorptions. Staining of non-related proteins (e.g. EBNA, at 72kD or VCA-p40 at 40kD) is not effected.

From the above described experiment it is apparent that immunostaining of viral VCA-p18 by human serum antibodies is specifically inhibited by preabsorption of these sera with BFRF3 fusion proteins, proving the immunochemical identity of VCA-p18 and the protein encoded by the BFRF3 reading frame. These experiments also indicate that reactivity of human sera with VCA-p18 is caused by interaction with a single viral protein.

Example 4:

20 Localization of immunoreactive epitopes by PEPSCAN.

Pentide synthesis and immunoscreening (PEPSCAN).

Peptides with a length of 12 amino acids (AA) and an overlap of 11 AA of the AA sequence of ORF BFRF3 and BDRF1 were synthesized by automated solid phase peptide synthesis onto chemically activated polyethylene pins as originally described by Geijsen et al. (Proc.Nat.Acad.Sci., USA, 83, 3998-4002, 1984). The immunoreactivity for EBV-specific antibodies, using 15 sera from healthy EBV seropositive donors was determined as described by Middeldorp and Melen (J.Virol.meth., 21, 147-159, 1988). The results of PEPSCAN analysis of 12 AA peptides derived from the VCA-p18 sequence are depicted in figure 4. The numbers on the X-axis represent the starting position on the VCA-p18 sequence of each 12-mer peptide analysed. Indicated on the Y-axis in figure 4 is the percentage of human sera reactive with a certain 12-mer peptide of the VCA-p18 protein encoded in the BFRF3 reading frame. Positive reaction is defined at 3 times the Standard Deviation above the mean reactivity of EBV-seronegative human serum on the same set of pins. From figure 4 three immunodominant domains can be defined:

Domain I: AA 120-140, Domain II: AA152-155, Domain III: AA 159-165 (These numbers again indicate the starting position of the 12-mer peptides used for PEPSCAN). Table 1 gives a detailed description of the most reactive peptides for each individual serum used in this study. In the first lane of Table 1 the number of the individual sera is indicated. The second lane indicates the PEPSCAN OD450 value (Optical Density at 450nm) for the most reactive peptide within domain I, which peptide is indicated in lane 3. Lane 4 and 5 represent similar data for domain II and III.

45

50

55

Table 1:

Immunological response of 15 EBV VCA-p18 positive sera.

(Only the response of the most reactive peptide located in the important epitope domains of VCA-p18 are illustrated).

Serum No.	Domain I (pept. 120-140)		Domain II (pept. 152-155)		Domain III (pept. 159-165)	
	Elisa OD ₄₅₀	Peptide A.A.-sequence	Elisa OD ₄₅₀	Peptide A.A.-sequence	Elisa OD ₄₅₀	Peptide A.A.-sequence
1	1.418	120-TAVAQSATPSVS-132	-	-	-	-
2	1.820	120-TAVAQSATPSVS-132	0.678	155-DTGSGGGGQPHD-167	0.423	162-GQPHDTAPRGAR-174
3	1.228	128-PSVSSSISSLRA-140	-	-	0.808	162-GQPHDTAPRGAR-174
4	1.230	128-PSVSSSISSLRA-140	-	-	0.761	162-GQPHDTAPRGAR-174
5	0.540	128-PSVSSSISSLRA-140	-	-	1.354	162-GQPHDTAPRGAR-174
6	0.731	129-SVSSSISSLRAA-141	-	-	1.441	162-GQPHDTAPRGAR-174
7	0.385	129-SVSSSISSLRAA-141	-	-	0.770	163-QPHDTAPRGAR-175
8	1.360	131-SSSISSLRAATS-143	-	-	1.343	160-GGGQPHDTAPRG-172
9	1.598	131-SSSISSLRAATS-143	0.510	153-AVDTGSGGGGQP-165	1.481	162-GQPHDTAPRGAR-174
10	1.591	131-SSSISSLRAATS-143	0.474	153-AVDTGSGGGGQP-165	1.481	162-GQPHDTAPRGAR-174
11	1.251	131-SSSISSLRAATS-143	0.958	152-AAVDTGSGGGGQ-164	0.774	162-GQPHDTAPRGAR-174
12	1.839	133-SISSLRAATSCA-145	-	-	0.407	162-GQPHDTAPRGAR-174
13	1.128	134-ISSLRAATSGAT-146	0.460	154-VDTGSGGGGQPH-166	1.535	162-GQPHDTAPRGAR-174
14	1.064	138-RAATSGATAAAS-150	-	-	1.319	162-GQPHDTAPRGAR-174
15	0.695	138-RAATSGATAAAS-150	-	-	0.644	162-GQPHDTAPRGAR-174

Example 5:

PEPSCAN analysis with rat and mouse monoclonal antibodies directed against BFRF3-encoded VCA-p18 protein, or the BdRF1-encoded VCA-p40 protein.

PEPSCAN analysis was performed, in a way similar to the procedure described in Example 4, for human sera, to delineate the position of linear epitopes detected by monoclonal antibodies.

Figures 5a and 5b show the PEPSCAN results of such an analysis using two rat monoclonal antibodies directed against VCA-p18 (EBV.OT15E and EBV.OT15I respectively), each detecting a different linear epitope.

Figure 6 shows the PEPSCAN results for a mouse monoclonal antibody (EBV.OT41A) directed against VCA-p40 encoded by the BdRF1 reading frame. Cross analysis of the two rat monoclonals on the VCA-p40 (BdRF1) sequence and vice-versa of the mouse monoclonal on the VCA-p18 (BFRF3) sequence gave clear negative results.

From figure 6 it is apparent that EBV.OT41A recognizes a distinct linear epitope in the C-terminal region of the VCA-p40 protein.

Example 6:

Selection of synthetic peptides derived from the BFRF3-encoded VCA-p18 protein by computer analysis and PEPSCAN, and analysis of immunoreactivity of these peptides with normal human donor sera.

Synthetic peptides were made by standard solid phase synthesis using t-BOC chemistry.

Peptides from BFRF3-encoded VCA-p18 protein were selected either on the basis of predicted high antigenicity using the computer program "antigenic index" developed by Jameson and Wolf (CABIOS 4, 181-186, 1988) [Peptides 1 and 2 in figure 7 were selected on this basis.] or on the basis of functional high antigen reactivity in PEPSCAN as described in Example 4. [Peptides 3 and 4 in figure 7 were selected on this basis, representing domain I plus II and domain III from table 1.] In addition, a combi-peptide was made (peptide 5 in figure 7) which represents a combination of the three most reactive domains identified by PEPSCAN, leaving out those peptide regions which show low PEPSCAN-reactivity.

Peptides 1-5 with amino acid sequences as indicated below were coated onto the solid phase, i.e. the wells of polystyrene microtiter plates, at 1 µg per ml in 0.05M NHCO₃ buffer at pH 9.6 overnight at 4°C. After washing twice with Phosphate-buffered saline (PBS) at pH7.4 the wells were filled with 100 µl of human serum, diluted 1:100 in PBS containing 0.05% Tween 20 (PBST) and incubated for 1 hour at 37°C.

After three PBS-T washes HRP labeled sheep anti-human IgG antibodies were added at the appropriate dilution in PBS-T and incubated for 1 hr at 37 °C. After three PBS-T washes, bound enzyme activity was detected using tetramethylbenzidine as substrate. The reaction was stopped at 30 minutes by adding 100 μ l 1M H₂SO₄. The absorbance was measured at 450 nm using a Multiscan photometer. Sera were tested for EBV-antibodies using standard immunofluorescence serology or immunoblot analysis as described before.

Peptides used were:

Peptide 1: H₂N-GVPRRQRAIDKRQRA-COOH

Peptide 2: H₂N-GQPHDTAPRGARKKQ-COOH

Peptide 3: H₂N-AVDTGSGGGGQPHDTAPRGARKKQ-COOH

Peptide 4: H₂N-STAVAQSATPSVSSSISLRAATSGATAAA-COOH

Peptide 5: Combi-peptide of peptide 4 and 3 linked by S-S-bridging, using extra cystein residues at the C-terminus of peptide 4 and the N-terminus of peptide 3.

Figure 7 shows the results of ELISA experiments using peptides 1-5 coated onto the solid phase and a random panel of sera from human healthy blood donors used at 1:100 dilution according to standard procedures.

Subsequently these sera were tested by immunoblot for the reactivity with viral VCA-p18.

From these experiments it is apparent that computer prediction based upon the "antigenic index" program have no predictive value with respect to the immunogenicity towards sera from naturally infected individuals as almost all sera were negative with peptide 1, while peptide 2 is reactive with only about 50% of the sera tested. Peptides selected on the basis of PEPSCAN show good reactivity with 60-80% of the sera tested respectively. Surprisingly the combi-peptide combining peptides 3 and 4 shows 95% reactivity with human sera. P18 immunoblot negative sera do not show any reactivity with the selected peptides.

Example 7:

Reactivity of combi-peptide 5, derived from the BFRF3-encoded VCA-p18 protein, with human serum antibodies of different subclasses.

The combi-peptide 5, with the amino acid sequence as described in Example 6, was used to analyse the reactivity of human serum immunoglobulins of different subclasses by means of ELISA (procedure as described in Example 6). In all cases combi-peptide 5 was used on the solid phase exactly as described in Example 6. Antibody reactivity was detected as described in Example 6 using 1:100 diluted sera. Results are shown in figure 8. IgM reactivity was detected in human sera pre-absorbed with Gull-Sorb (Gull Laboratories Inc., Salt Lake City, Utah, USA) according to the instructions of the manufacturer, in order to inhibit IgG-reactivity. IgM antibodies were detected with HRP-labeled sheep anti-human IgM antibodies specific for the heavy chain of human IgM.

IgA reactivity was also detected in Gull Sorb treated sera using anti-IgA specific HRP labeled second antibodies.

Sera for figure 8a (IgG) were obtained from 76 healthy seropositive blood donors, positive for VCA-IgG by standard serology and 9 donors negative for EBV antibodies.

Sera for figure 8b (IgM) were obtained from 26 mononucleosis patients positive for VCA-IgM by standard serology and 18 healthy donors negative for VCA-IgM but positive for VCA-IgG.

Sera for figure 8c (IgA) were obtained from 35 confirmed Nasopharyngeal Carcinoma patients from whom no IgA-specific data were available, but all of which were IgG-VCA positive and from 7 healthy VCA-IgG positive donors.

From the above experiments it can be seen that the VCA-p18 derived combi peptide can be used for specific detection of IgG, IgM and IgA antibodies to EBV-VCA, with sensitivity similar or better than standard serological techniques. EBV-negative sera are negative in all cases.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- 5 (i) APPLICANT:
(A) NAME: Akzo N.V.
(B) STREET: Velperweg 76
(C) CITY: Arnhem
(E) COUNTRY: The Netherlands
(F) POSTAL CODE (ZIP): 6824 BM
- 10 (ii) TITLE OF INVENTION: Peptides and nucleic acid sequences
related to the Epstein-Barr virus.
- (iii) NUMBER OF SEQUENCES: 22
- 15 (iv) COMPUTER READABLE FORM:
(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: Patentin Release #1.0, Version #1.25
(EPO)

20 (2) INFORMATION FOR SEQ ID NO: 1:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 538 base pairs
(B) TYPE: nucleic acid
25 (C) STRANDEDNESS: double
(D) TOPOLOGY: unknown
- (ii) MOLECULE TYPE: DNA (genomic)
- (vi) ORIGINAL SOURCE:
(A) ORGANISM: Epstein-Barr virus
- 30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

CATGATGGCA CGCCGGCTGC CCAAGCCCAC CCTCCAGGGG AGGCTGGAGG CGGATTTTCC 60

35 AGACAGTCCC CTGCTTCCTA AATTTCAAGA GCTGAACCAG AATAATCTCC CCAATGATGT 120

TTTTCGGGAG GCTCAAAGAA GTTACCTGGT ATTTCTGACA TCCCAGTTCT GCTACGAAGA 180

GTACGTGCAG AGGACTTTTG GGGTGCCTCG GCGCCAACGC GCCATAGACA AGAGGCAGAG 240

40 AGCCAGTGTG GCTGGGGCTG GTGCTCATGC ACACCTTGGC GGGTCATCCG CCACCCCGT 300

CCAGCAGGCT CAGGCCGCCG CATCCGCTGG GACCGGGGCC TTGGCATCAT CAGCGCCGTC 360

CACGGCCGTA GCCCAGTCCG CGACCCCTC TGTTCCTTCA TCTATTAGCA GCCTCCGGGC 420

45 CGCGACTTCG GGGGCGACTG CCGCCGCCCTC CGCCGCCGCA GCCGTCGATA CCGGGTCAGG 480

TGGCGGGGGA CAACCCACG ACACCGCCCC ACGCGGGGCA CGTAAGAAAC AGTAGCCC 538

50

55

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 176 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Epstein-Barr virus

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Met Ala Arg Arg Leu Pro Lys Pro Thr Leu Gln Gly Arg Leu Glu Ala
 1 5 10 15
 Asp Phe Pro Asp Ser Pro Leu Leu Pro Lys Phe Gln Glu Leu Asn Gln
 20 25 30
 Asn Asn Leu Pro Asn Asp Val Phe Arg Glu Ala Gln Arg Ser Tyr Leu
 35 40 45
 Val Phe Leu Thr Ser Gln Phe Cys Tyr Glu Glu Tyr Val Gln Arg Thr
 50 55 60
 Phe Gly Val Pro Arg Arg Gln Arg Ala Ile Asp Lys Arg Gln Arg Ala
 65 70 75 80
 Ser Val Ala Gly Ala Gly Ala His Ala His Leu Gly Gly Ser Ser Ala
 85 90 95
 Thr Pro Val Gln Gln Ala Gln Ala Ala Ala Ser Ala Gly Thr Gly Ala
 100 105 110
 Leu Ala Ser Ser Ala Pro Ser Thr Ala Val Ala Gln Ser Ala Thr Pro
 115 120 125
 Ser Val Ser Ser Ser Ile Ser Ser Leu Arg Ala Ala Thr Ser Gly Ala
 130 135 140
 Thr Ala Ala Ala Ser Ala Ala Ala Ala Val Asp Thr Gly Ser Gly Gly
 145 150 155 160
 Gly Gly Gln Pro His Asp Thr Ala Pro Arg Gly Ala Arg Lys Lys Gln
 165 170 175

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1038 base pairs
 (B) TYPE: nucleic acid

(C) STRANDEDNESS: double
(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Epstein-Barr virus

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

ATGCTATCAG GTAACGCAGG AGAAGGAGCA ACAGCCTGCG GAGGTTCTGGC CGCCGCGGGC 60
CAGGACCTCA TCAGCGTCCC CCGCAACACC TTTATGACAC TGCTTCAGAC CAACCTGGAC 120
AACAAACCGC CGAGGCAGAC CCCGCTACCC TACGCGGCCC CGCTGCCCCC CTTTTCACAC 180
CAGGCAATAG CCACCGCGCC TTCCTACGGT CCTGGGGCCG GAGCGGTCTGC CCCGGCCGGC 240
GGCTACTTTA CCTCCCCAGG AGGTTACTAC GCCGGGCCCC CGGGCGGGGA CCCGGGTGCC 300
TTCTTGGCGA TGGACGCTCA CACCTACCAC CCCCACCCAC ACCCCCCTCC GGCCTACTTT 360
GGCTTGCCGG GCCTCTTTGG CCCCCCTCCA CCGTGCTCTC CTTACTACGG ATCCCACTTG 420
CGGGCAGACT ACGTCCCCGC TCCCTCGCGA TCCAACAAGC GGAAAAGAGA CCCCAGGAG 480
GATGAAGAAG GCGGGGGGCT ATTCCCGGGG GAGGACGCCA CCCTCTACCG CAAGGACATA 540
GCGGGCCTCT CCAAGAGTGT GAATGAGTTA CAGCACACGC TACAGGCCCT GCGCCGGGAG 600
ACGCTGTCCT ACGGCCACAC CGGAGTCGGA TACTGCCCCC AGCAGGGCCC CTGCTACACC 660
CACTCGGGGC CTTACGGATT TCAGCCTCAT CAAAGCTACG AAGTGCCAG ATACGTCCCT 720
CATCCGCCCC CACCACCAAC TTCTCACCAG GCAGCTCAGG CGCAGCCTCC ACCCCCGGGC 780
ACACAGGCCC CCGAAGCCCA CTGTGTGGCC GAGTCCACGA TCCCTGAGGC GGGAGCAGCC 840
GGGAACTCTG GACCCCGGGA GGACACCAAC CCTCAGCAGC CCACCACCGA GGGCCACCAC 900
CGCGGAAAGA AACTGGTGCA GGCCTCTGCG TCCGGAGTGG CTCAGTCTAA GGAGCCCACC 960
ACCCCAAGG CCAAGTCTGT GTCAGCCAC CTCAAGTCCA TCTTTTGC GAATTGCTG 1020
AATAAACGCG TGGCTTGA 1038

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 345 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Epstein-Barr virus

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Met Leu Ser Gly Asn Ala Gly Glu Gly Ala Thr Ala Cys Gly Gly Ser
 1 5 10 15
 Ala Ala Ala Gly Gln Asp Leu Ile Ser Val Pro Arg Asn Thr Phe Met
 20 25 30
 Thr Leu Leu Gln Thr Asn Leu Asp Asn Lys Pro Pro Arg Gln Thr Pro
 35 40 45
 Leu Pro Tyr Ala Ala Pro Leu Pro Pro Phe Ser His Gln Ala Ile Ala
 50 55 60
 Thr Ala Pro Ser Tyr Gly Pro Gly Ala Gly Val Ala Pro Ala Gly
 65 70 75 80
 Gly Tyr Phe Thr Ser Pro Gly Gly Tyr Tyr Ala Gly Pro Ala Gly Gly
 85 90 95
 Asp Pro Gly Ala Phe Leu Ala Met Asp Ala His Thr Tyr His Pro His
 100 105 110
 Pro His Pro Pro Pro Ala Tyr Phe Gly Leu Pro Gly Leu Phe Gly Pro
 115 120 125
 Pro Pro Pro Val Pro Pro Tyr Tyr Gly Ser His Leu Arg Ala Asp Tyr
 130 135 140
 Val Pro Ala Pro Ser Arg Ser Asn Lys Arg Lys Arg Asp Pro Glu Glu
 145 150 155 160
 Asp Glu Glu Gly Gly Gly Leu Phe Pro Gly Glu Asp Ala Thr Leu Tyr
 165 170 175
 Arg Lys Asp Ile Ala Gly Leu Ser Lys Ser Val Asn Glu Leu Gln His
 180 185 190
 Thr Leu Gln Ala Leu Arg Arg Glu Thr Leu Ser Tyr Gly His Thr Gly
 195 200 205
 Val Gly Tyr Cys Pro Gln Gln Gly Pro Cys Tyr Thr His Ser Gly Pro
 210 215 220
 Tyr Gly Phe Gln Pro His Gln Ser Tyr Glu Val Pro Arg Tyr Val Pro
 225 230 235 240
 His Pro Pro Pro Pro Pro Thr Ser His Gln Ala Ala Gln Ala Gln Pro
 245 250 255
 Pro Pro Pro Gly Thr Gln Ala Pro Glu Ala His Cys Val Ala Glu Ser

260 265 270
 Thr Ile Pro Glu Ala Gly Ala Ala Gly Asn Ser Gly Pro Arg Glu Asp
 275 280 285
 5 Thr Asn Pro Gln Gln Pro Thr Thr Glu Gly His His Arg Gly Lys Lys
 290 295 300
 Leu Val Gln Ala Ser Ala Ser Gly Val Ala Gln Ser Lys Glu Pro Thr
 10 305 310 315 320
 Thr Pro Lys Ala Lys Ser Val Ser Ala His Leu Lys Ser Ile Phe Cys
 325 330 335
 Glu Glu Leu Leu Asn Lys Arg Val Ala
 15 340 345

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Epstein-Barr virus

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

Ala Val Asp Thr Gly Ser Gly Gly Gly Gln Pro His Asp Thr Ala
 5 10 15
 Pro Arg Gly Ala Arg Lys Lys Gln
 20

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Epstein-Barr virus

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

Ser Thr Ala Val Ala Gln Ser Ala Thr Pro Ser Val Ser Ser Ser Ile
5 10 15

Ser Ser Leu Arg Ala Ala Thr Ser Gly Ala Thr Ala Ala Ala
20 25

(2) INFORMATION FOR SEQ ID NO: 7:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 15 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Epstein-Barr virus

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

Gly Val Pro Arg Arg Gln Arg Ala Ile Asp Lys Arg Gln Arg Ala
5 10 15

(2) INFORMATION FOR SEQ ID NO: 8:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 15 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Epstein-Barr virus

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

Gly Gln Pro His Asp Thr Ala Pro Arg Gly Ala Arg Lys Lys Gln
5 10 15

(2) INFORMATION FOR SEQ ID NO: 9:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Epstein-Barr virus

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

Thr Ala Val Ala Gln Ser Ala Thr Pro Ser Val Ser
5 10

(2) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Epstein-Barr virus

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

Pro Ser Val Ser Ser Ser Ile Ser Ser Leu Arg Ala
5 10

(2) INFORMATION FOR SEQ ID NO: 11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Epstein-Barr virus

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

Ser Val Ser Ser Ser Ile Ser Ser Leu Arg Ala Ala
5 10

(2) INFORMATION FOR SEQ ID NO: 12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Epstein-Barr virus

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

Ser Ser Ser Ile Ser Ser Leu Arg Ala Ala Thr Ser
5 10

(2) INFORMATION FOR SEQ ID NO: 13:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Epstein-Barr virus

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

Ser Ile Ser Ser Leu Arg Ala Ala Thr Ser Gly Ala
5 10

(2) INFORMATION FOR SEQ ID NO: 14:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Epstein-Barr virus

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

Ile Ser Ser Leu Arg Ala Ala Thr Ser Gly Ala Thr
5 10

(2) INFORMATION FOR SEQ ID NO: 15

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Epstein-Barr virus

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15

Arg Ala Ala Thr Ser Gly Ala Thr Ala Ala Ala Ser
5 10

(2) INFORMATION FOR SEQ ID NO: 16

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Epstein-Barr virus

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

Ala Ala Val Asp Thr Gly Ser Gly Gly Gly Gly Gln
5 10

(2) INFORMATION FOR SEQ ID NO: 17

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Epstein-Barr virus

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

Ala Val Asp Thr Gly Ser Gly Gly Gly Gly Gln Pro
5 10

(2) INFORMATION FOR SEQ ID NO: 18

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Epstein-Barr virus

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

Val Asp Thr Gly Ser Gly Gly Gly Gly Gln Pro His
5 10

(2) INFORMATION FOR SEQ ID NO: 19

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Epstein-Barr virus

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

Asp Thr Gly Ser Gly Gly Gly Gly Gln Pro His Asp
5 10

(2) INFORMATION FOR SEQ ID NO: 20:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Epstein-Barr virus

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

Gly Gly Gly Gln Pro His Asp Thr Ala Pro Arg Gly
5 10

(2) INFORMATION FOR SEQ ID NO: 21:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Epstein-Barr virus

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:

Gly Gln Pro His Asp Thr Ala Pro Arg Gly Ala Arg
5 10

(2) INFORMATION FOR SEQ ID NO: 22:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 12 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Epstein-Barr virus

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:

Gln Pro His Asp Thr Ala Pro Arg Gly Ala Arg Lys
5 10

35 Claims

1. Peptide, immunochemically reactive with antibodies to the Epstein Barr Virus, comprising at least part of the VCA-p18 or VCA-p40 protein, encoded within the EBV open reading frames BFRF3 and BpRF1 respectively, or a functional variant thereof.
2. Peptide according to claim 1, comprising at least part of the amino acid sequence shown in SEQ ID No.: 2 or a functional variant thereof.
3. Peptide according to claim 1, comprising at least part of the amino acid sequence shown in SEQ ID No.: 4 or a functional variant thereof.
4. Peptide according to claim 2, comprising at least one of the amino acid sequences chosen from the amino acid sequences as shown in SEQ ID NO:5 and SEQ ID NO:6.
5. Peptide according to claim 4, comprising the amino acid sequence as shown in SEQ ID NO:5 linked to the amino acid sequences as shown in SEQ ID NO:6.
6. Nucleic acid sequence encoding a peptide according to any of claims 1-5.
7. Nucleic acid sequence, comprising at least part of the nucleic acid sequence as shown in SEQ ID NO: 1.

8. Nucleic acid sequence, comprising at least part of the nucleic acid sequence as shown in SEQ ID NO.: 3.
9. A recombinant vector molecule comprising a nucleic acid sequence according to any of claims 6-8.
10. A host cell transformed or transfected with a recombinant vector according to claim 9.
11. Antibody directed to a peptide according to any of claims 1-5.
12. Antibody according to claim 11, the antibody being a monoclonal antibody.
13. Monoclonal antibody, having the same reactivity with VCA-p18 as monoclonal antibody EBV.OT15E or EBV.OT15I produced by the rat-mouse hybridoma cell lines deposited with the European Collection of Animal Cell Cultures (ECACC), Porton Down (UK), under deposit No. 93020413 and 93020412 respectively.
14. Monoclonal antibody having the same reactivity with VCA-p40 as monoclonal antibody EBV.OT41A produced by the mouse-mouse hybridoma cell line deposited with the European Collection of Animal Cell Cultures (ECACC), Porton Down (UK), under deposit No. 93020414.
15. Immortalized cell line capable of producing monoclonal antibodies according to claim 13 or 14.
16. Immortalized cell line according to claim 15, deposited with the European Collection of Animal Cell Cultures (ECACC), Porton Down (UK), under deposit No. 93020413 or 93020412 or 93020414.
17. Anti-idiotypic antibody reactive with the antibody according to claim 11.
18. Immunochemical reagent comprising one or more peptide(s) according to any of claims 1-5.
19. Immunological reagent comprising one or more antibodies according to any of claims 11-14.
20. Method for the detection of Epstein-Barr virus in a sample characterized in that an antibody according to any of claims 11-14 is brought into contact with a sample whereafter the presence of immune complexes formed is detected which is a measure for the presence of Epstein-Barr virus in the sample.
21. Method for the detection of antibodies directed against Epstein-Barr Virus in a sample, characterized in that an immunochemical reagent according to claim 14 is brought into contact with the sample and the presence of immune complexes formed between the peptide and antibodies in the sample is detected, which is a measure for the presence of Epstein Barr Virus antibodies in the sample.
22. Method for the detection of Epstein-Barr Virus in a sample characterized in that one or more peptides according to any of claims 1-5 are brought into contact with sample and antibodies directed Epstein-Barr Virus, whereafter the presence of immune complexes formed is detected and from this the presence of Epstein-Barr Virus in the sample is determined.
23. Method for the amplification and the detection of an Epstein-Barr Virus nucleic acid sequence in a sample using at least one nucleic acid sequence or fragment thereof according to claim 6-8 as primer(s) in order to perform a nucleic acid amplification of said Epstein-Barr Virus nucleic acid sequence and to detect the amplified sequence.
24. Test kit for carrying out a method according to any of claims 20-22.
25. Test amplification kit for carrying out an amplification method according to claim 23.

FIGURE 1




FIGURE 2




FIGURE 3

IMMUNO-ADSORPTION





FIGURE 4

VCA-p18 EPIOTOPE RECOGNITION of 15 EBV+ human sera



Series 1

FIGURE 5A
PEPSCAN of BFRF3
EBV.OT15E




FIGURE 5B
PEPSCAN of BFRF3
EBV.OT15I




FIGURE 6




FIGURE 7

VCA-p18 peptides Reactivity of 43 healthy blood donors




FIGURE 8A

VCA-p18 peptide ELISA (IgG)
(76 VCA-I.F. positive sera)





FIGURE 8B

EBV-IgM: VCA-p18 peptide
Reactivity of 18 IgM-negative and 26
IgM-positive sera.



111

111



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number: **0 574 048 A3**

EUROPEAN PATENT APPLICATION

Application number: **93200691.9**

Date of filing: **10.03.93**

Int. Cl.⁵: **C12N 15/38, C12P 21/08,
C12Q 1/68, C07K 13/00,
C07K 15/28, G01N 33/536,
G01N 33/68**

Priority: **13.03.92 EP 92200721**

Date of publication of application:
15.12.93 Bulletin 93/50

Designated Contracting States:
**AT BE CH DE DK ES FR GB GR IE IT LI LU MC
NL PT SE**

Date of deferred publication of the search report:
02.03.94 Bulletin 94/09

Applicant: **AKZO N.V.
Velperweg 76
NL-6824 BM Arnhem(NL)**


Inventor: **Middeldorp, J.M.
Staringstraat 151
NL-5343 GD Oss(NL)**

Representative: **Hermans, Franciscus G.M. et
al
P.O. Box 20
NL-5340 BH Oss (NL)**

Peptides and nucleic acid sequences related to Epstein Barr Virus.

The present invention relates to peptides immunochemically reactive with antibodies to the Epstein-Barr virus (EBV), comprising at least part of the VCA-p18 or VCA-p40 protein, encoded within the EBV open reading frames BFRF3 and BdRF1 respectively, or a functional variant thereof. The invention further relates to nucleic acid sequences encoding these peptides, monoclonal antibodies against these peptides, cell lines capable of producing monoclonal antibodies and anti-idiotypic antibodies. The invention also relates to recombinant vector molecules comprising a nucleic acid sequence according to the invention and host cells transformed or transfected with these vector molecules. The invention is further concerned with immunological reagents and methods for the detection of EBV or anti-EBV antibodies and a method for the amplification and detection of Epstein Barr viral nucleic acid.

FIGURE 1





European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 93 20 0691

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 5)
D, Y	NATURE vol. 310, 19 July 1984, MACMILLAN JOURNALS LTD., LONDON, GB; pages 207 - 211 R. BAER ET AL. 'DNA sequence and expression of the B95-8 Epstein-Barr virus genome' * page 209, left column, line 14; table 1; page 209, left column, line 75; table 1 *	1, 2, 4-7 , 9-13, 15-20, 22-25	C 12 N 15/38 C 12 P 21/08 C 12 Q 1/68 C 07 K 13/00 C 07 K 15/28 G 01 N 33/536 G 01 N 33/68
Y	J. GEN. VIROL. vol. 56, 1981, SOC. GEN. MICROBIOL., LONDON, GB; pages 105 - 118 G.J. BAYLISS AND H. WOLF 'The regulated expression of Epstein-Barr virus. III. Proteins specified by EBV during the lytic cycle' * page 115, line 9 - page 117, line 5; figures 1-6; table 1 *	1, 2, 4-7 , 9-13, 15-20, 22-25	
Y	WO-A-9 109 127 (SCHERING CORPORATION) 27 June 1991 * page 2, line 2 - page 3, line 6; page 8, line 5 - page 10, line 33; claims 1-12 *	1, 2, 4-7 , 9-13, 15-20, 22-25	TECHNICAL FIELDS SEARCHED (Int. Cl. 5)
Y	WO-A-9 108 224 (THE COUNCIL OF THE QUEENSLAND INSTITUTE OF MEDICAL RESEARCH) 13 June 1991 * the whole document *	1, 2, 4-7 , 9-13, 15-20, 22-25	C 07 K C 12 P G 01 N
Y	EP-A-0 316 170 (G GENE GALWAY LIMITED) 17 May 1989 * the whole document *	1, 2, 4-7 , 9-13, 15-20, 22-25	
Y	WO-A-9 102 091 (NORTHWESTERN UNIVERSITY) 21 February 1991 * the whole document *	23, 25	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 02-06-1993	Examiner HORNIG H
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	



European Patent
Office

CLAIMS INCURRING FEES

The present European patent application comprised at the time of filing more than ten claims.

☐

All claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for all claims.

☐

Only part of the claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims and for those claims for which claims fees have been paid,

namely claims:

☐

No claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims.

LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirement of unity of invention and relates to several inventions or groups of inventions.

namely:

see sheet -B-

☐

All further search fees have been paid within the fixed time limit. The present European search report has been drawn up for all claims.

☐

Only part of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the inventions in respect of which search fees have been paid.

namely claims:

☒

None of the further search fees has been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the invention first mentioned in the claims.

namely claims

2,4,5,7,13(completely)1,6,9-12,15-20,22-25 (partially)



European Patent
Office

Sheet 8

EP 93200691

LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirement of unity of invention and relates to several inventions or groups of inventions, namely:

1. Claims 2,4,5,7,13 (completely) and claims 1,6,9-12,15-20,22-25 (partially)
Nucleic acid sequence as a fragment thereof coding for VCA-P18 or a fragment thereof, which is immuno-chemically reactive with EBV antibodies; nucleic acid sequence BFRF3 or a fragment thereof coding for said protein or fragment thereof; immuno-chemical reagent comprising said peptide, fragment or polypeptide. Monoclonal antibody against said polypeptide or fragment thereof; immortalized cell line producing said antibody; anti-idiotypic antibody reactive with said antibody; method for detecting of antibodies using said VCA-P18 or fragments thereof; method for the amplification and the detection of an EBV nucleic acid sequence using said BFRF3-DNA sequence information; test kit for carrying out a nucleic acid amplification using said polypeptide or said nucleic acid sequence.

2. Claims 3,8,14,21 (completely) and claims 1,6,9-12,15-20,22-25 (partially)
Nucleic acid sequence or a fragment thereof coding for VCA-P40 or a fragment thereof, which is immuno-chemically reactive with EBV antibodies; nucleic sequence BdRF1 or a fragment thereof coding for said protein or fragment thereof; immuno-chemical reagent comprising said peptide, fragment or polypeptide; monoclonal antibody against said polypeptide or fragment thereof; immortalized cell line producing said antibody; anti-idiotypic antibody reactive with said antibody; method for detecting of antibodies using said VCA-P40 or fragment thereof; method for the amplification and the detection of an EBV-nucleic acid sequence using said BdRF1-DNA sequence information; test kit for carrying out an immuno-assay or test amplification kit for carrying out a nucleic acid amplification using said polypeptide or said nucleic acid sequence.